

## Neutrino Mass, Inflation, and Dark Energy with the CMB

Prof. Neelima Sehgal Stony Brook University

DOE Cosmic Visions Dark Energy Oct. 1st, 2015

• Intro to ACTPol, AdvACT, CMB-S4

- Intro to ACTPol, AdvACT, CMB-S4
- Neutrino Mass from CMB Lensing

- Intro to ACTPol, AdvACT, CMB-S4
- Neutrino Mass from CMB Lensing
- Primordial Gravity Waves and Delensing


- Intro to ACTPol, AdvACT, CMB-S4
- Neutrino Mass from CMB Lensing
- Primordial Gravity Waves and Delensing
- CMB Halo Lensing/CMB Lensing Cross-Correlations as Probes of Dark Energy

- Intro to ACTPol, AdvACT, CMB-S4
- Neutrino Mass from CMB Lensing
- Primordial Gravity Waves and Delensing
- CMB Halo Lensing/CMB Lensing Cross-Correlations as Probes of Dark Energy
- Conclusion

- Intro to ACTPol, AdvACT, CMB-S4
- Neutrino Mass from CMB Lensing
- Primordial Gravity Waves and Delensing
- CMB Halo Lensing/CMB Lensing Cross-Correlations as Probes of Dark Energy
- Conclusion

#### ACTPol / AdvACT

#### ACTPol / AdvACT



#### ACTPol / AdvACT



ACTpol - Observes from 2013 - 2015

ACTpol - Observes from 2013 - 2015 4000 sq deg (10% of sky)

```
ACTpol - Observes from 2013 - 2015
4000 sq deg (10% of sky)
4 x better sensitivity than Planck
```


```
ACTpol - Observes from 2013 - 2015
4000 sq deg (10% of sky)
4 x better sensitivity than Planck
5 x better resolution than Planck
```

ACTpol - Observes from 2013 - 2015 4000 sq deg (10% of sky)

4 x better sensitivity than Planck

5 x better resolution than Planck

AdvACTpol - 2016 - 2018




ACTpol - Observes from 2013 - 2015 4000 sq deg (10% of sky)

4 x better sensitivity than Planck

5 x better resolution than Planck

AdvACTpol - 2016 - 2018 20,000 sq deg (50% of sky)




ACTpol - Observes from 2013 - 2015 4000 sq deg (10% of sky)

4 x better sensitivity than Planck

5 x better resolution than Planck

AdvACTpol - 2016 - 2018 20,000 sq deg (50% of sky) 4 x better sensitivity than Planck



ACTpol - Observes from 2013 - 2015 4000 sq deg (10% of sky)

4 x better sensitivity than Planck


5 x better resolution than Planck

AdvACTpol - 2016 - 2018

20,000 sq deg (50% of sky)

4 x better sensitivity than Planck

5 x better resolution than Planck



ACTpol - Observes from 2013 - 2015 4000 sq deg (10% of sky)

4 x better sensitivity than Planck


5 x better resolution than Planck

AdvACTpol - 2016 - 2018


20,000 sq deg (50% of sky)


4 x better sensitivity than Planck

5 x better resolution than Planck



AdvACT funded by NSF MSIP in June 2015





#### **Building for Discovery**

Strategic Plan for U.S. Particle Physics in the Global Context

Report of the Particle Physics Project Prioritization Panel (P5)

	Scenarios				Science Drivers					
Project/Activity	Scenario A	Scenario B	Scenario C	Higgs	Neutrinos	Dark Matter	Cosm. Accel.	The Unknown	Technique (Frontier)	
Large Projects										
Muon program: Mu2e, Muon g-2	Y, Mu2e small reprofile needed	Y	Y					~	ı	
HL-LHC	Y	Y	Y	~		1		~	Ε	
LBNF + PIP-II	Y, delayed relative to Scenario B.	Υ	Y, enhanced		1			1	I,C	
ILC	R&D only	R&D, possibly small handware contri- butions. See text.	Y	~		1		1	Ε	
NuSTORM	N	N	N		~				ı	
RADAR	N	N	N		~				ı	
Medium Projects										
LSST	Y	Y	Y		/		1		С	
DM G2	Y	Y	Y			1			С	
Small Projects Portfolio	Y	Υ	Y		~	~	~	~	All	
Accelerator R&D and Test Facilities	Y, reduced	Y, redirection to PIP-II development	Y, enhanced	~	~	~		~	E,I	
CMB-S4	Y	Y	Y		~		1		С	

#### **Building for Discovery**


Strategic Plan for U.S. Particle Physics in the Global Context

Report of the Particle Physics Project Prioritization Panel (P5)


	Scenarios			Science Drivers					
Project/Activity	Scenario A	Scenario B	Scenario C	Higgs	Neutrinos	Dark Matter	Cosm. Accel.	The Unknown	Technique (Frontier)
Large Projects									
Muon program: Mu2e, Muon g-2	Y, Mu2e small reprofile	Y	Y					~	1
HL-LHC	Υ	Υ	Y	~		1		1	Е
LBNF + PIP-II	Y, delayed relative to Scenario B.	Υ	Y, enhanced		/			1	I,C
ILC	R&D only	R&D, possibly small hardware contri- butions. See text.	Υ	~		1		1	Ε
NuSTORM	N	N	N		~				1
RADAR	N	N	N		~				1
Medium Projects									
LSST	Υ	Y	Υ		~		1		С
DM G2	Y	Y	Y			~			С
Small Projects Portfolio	Υ	Y	Υ		~	~	~	1	All
Accelerator R&D and Test Facilities	Y, reduced	y redirection to	Y, enhanced	1	~	~		~	E,I
CMB-S4	Y	Y	Y		~		~		С

- Intro to ACTPol, AdvACT, CMB-S4
- Neutrino Mass from CMB Lensing
- Primordial Gravity Waves and Delensing
- CMB Halo Lensing/CMB Lensing Cross-Correlations as Probes of Dark Energy
- Conclusion


#### Unlensed CMB



#### Lensed CMB




# Measurements of CMB Lensing on Large Scales



Planck Paper 15, 2015 (1502.01591)

# Measurements of CMB Lensing on Large Scales



Planck Paper 15, 2015 (1502.01591)

Blanchard & Schneider 1987 (first idea of detectability)

Zaldarriaga & Seljak 1997 (first lensing estimators)

Hu 2001 Hu & Okamoto 2002 (optimal lensing estimators)

Smith, Zahn, Dore 2007 (first indirect detection)

Das et al. 2011 - ACT (first direct detection)

van Engelen et al. 2012 - SPT (second direct detection)

Planck Collaboration 2013 (detection with S/N = 25)

Planck Collaboration 2015 (detection with S/N = 40)

# CMB Lensing Power Spectrum Sensitive to Neutrino Mass








Figure credit: A. van Engelen











Data Set	$f_{sky}$	Map noise (μK-arcmin)	$\sigma(\sum m_{\nu}) \text{ (eV)}$	$\sigma(\sum m_{\nu}) \text{ (eV)}$
		at 150 GHz	CMB alone	with BAO
Planck	0.8	43	0.20	0.12
ACTPol	0.1	20	0.09	0.06
AdvACT	0.5	7	0.06	0.04

#### **CMB-S4** Forecasts



Allison et al., 1509.07471, 2015

#### **CMB-S4** Forecasts




Allison et al., 1509.07471, 2015


#### Outline


- Intro to ACTPol, AdvACT, CMB-S4
- Neutrino Mass from CMB Lensing
- Primordial Gravity Waves and Delensing
- CMB Halo Lensing/CMB Lensing Cross-Correlations as Probes of Dark Energy
- Conclusion

$$T^{L}(\hat{n}) = T^{U}(\hat{n} + \nabla \phi(\hat{n}))$$


$$T^L(\hat{n}) = T^U(\hat{n} + \underline{\nabla \phi(\hat{n})})$$
 Deflection angle








 Delensing = undo the lensing of the primordial CMB due to Large-scale structure (LSS)



Shift pixels backward using LSS map to reconstruct unlensed CMB


 Delensing = undo the lensing of the primordial CMB due to Large-scale structure (LSS)



Shift pixels backward using LSS map to reconstruct unlensed CMB

 Need template of LSS from either internal CMB lens map reconstruction and/or LSS tracer like CIB (see e.g. Sherwin & Schmittfull, 2015, 1502.05356)


Want to measure the amplitude (r) and scale dependence (n<sub>T</sub>) of primordial B-mode power



Simard, Hanson, Holder 2014, 1410.0691

Want to measure the amplitude (r) and scale dependence (n<sub>T</sub>) of primordial B-mode power

 $n_T = -r/8$  is consistency relation of single-field, slow roll inflation




Simard, Hanson, Holder 2014, 1410.0691

Want to measure the amplitude (r) and scale dependence (n<sub>T</sub>) of primordial B-mode power

 $n_T = -r/8$  is consistency relation of single-field, slow roll inflation

B-modes from gravitational lensing are a contaminant



Simard, Hanson, Holder 2014, 1410.0691

Error on r proportional to B-mode lens power

Error on r proportional to B-mode lens power

$$\sigma(r) \propto \sum_{l} \sqrt{\frac{2}{(2l+1)f_{\rm sky}}} (C_l^{\rm BB,lens} + N_l^{\rm BB})$$

Error on r proportional to B-mode lens power


$$\sigma(r) \propto \sum_{l} \sqrt{\frac{2}{(2l+1)f_{\rm sky}}} (C_l^{\rm BB,lens} + N_l^{\rm BB})$$

 With map of LSS and CMB E-mode map, can estimate CMB B-mode map from lensing


Error on r proportional to B-mode lens power

$$\sigma(r) \propto \sum_{l} \sqrt{\frac{2}{(2l+1)f_{\rm sky}}} (C_l^{\rm BB,lens} + N_l^{\rm BB})$$

- With map of LSS and CMB E-mode map, can estimate CMB B-mode map from lensing
  - Need range of scales (100 < 1 < 1000), including small-scales as they contribute to large-scale
     B-mode power




AdvACT will survey 50% of sky



AdvACT will survey 50% of sky


have 5 frequency channels



AdvACT will survey 50% of sky

have 5 frequency channels

have a HWP to get to ell ~ 50




AdvACT will survey 50% of sky


have 5 frequency channels

have a HWP to get to ell ~ 50

have small-scale CMB to delens



#### CMB-S4: Inflation Forecast



#### Outline

- Intro to ACTPol, AdvACT, CMB-S4
- Neutrino Mass from CMB Lensing
- Primordial Gravity Waves and Delensing
- CMB Halo Lensing/CMB Lensing Cross-Correlations as Probes of Dark Energy
- Conclusion


#### CMB Lensing on Small Scales

#### Unlensed CMB

Noiseless unlensed CMB sim

20' x 20' patch

Mostly gradient



#### CMB Lensing on Small Scales


#### Lensed CMB

Noiseless lensed CMB sim

20' x 20' patch

Lensed by  $M_{180} = 2 \times 10^{15} \text{ Msun}$ 

Lensing signal on Mpc / arcmin scales



#### CMB Lensing on Small Scales


Difference of lensed and unlensed CMB

20' x 20' patch


Characteristic dipole along the direction of gradient

Dipole signal is of the order of ~I-I0uK

#### Difference Map



# ACTPol First Season Maps



# Stacked ~12,000 CMASS Galaxies

# Stacked ~12,000 CMASS Galaxies

Galaxies from SDSS-III/BOSS DR10

CMASS ("constant mass")
galaxies have similar
selection as LRGs


# Stacked ~12,000 CMASS Galaxies

Galaxies from SDSS-III/BOSS DR10

CMASS ("constant mass")
galaxies have similar
selection as LRGs

Galaxies chosen because have optical weak lensing mass estimates


#### Weak lensing using CFHTLS



$$M_{200} = (2.3 \pm 0.1) \times 10^{13} h^{-1} M_{\odot}$$

# ACTPol: First Detection of Lensing of the CMB by Dark Matter Halos


Madhavacheril, Sehgal, et. al., PRL, 114, 2015




We detect halo lensing from 12,000 stacked CMASS galaxies at S/N of 3.2 sigma

Best fit:  $M_{200} = (2.0 \pm 0.7) \times 10^{13} h^{-1} M_{\odot}$  and  $c_{200} = 5.4 \pm 0.8$ 


#### AdvACT: Dark Energy Forecast



### AdvACT: Dark Energy Forecast




#### CMB Lens Cross-Correlations



Das et al, 2013, "Can CMB Lensing Help Cosmic Shear Surveys"

#### CMB Lens Cross-Correlations



Das et al, 2013, "Can CMB Lensing Help Cosmic Shear Surveys"

#### Outline

- Intro to ACTPol, AdvACT, CMB-S4
- Neutrino Mass from CMB Lensing
- Primordial Gravity Waves and Delensing
- CMB Halo Lensing/CMB Lensing Cross-Correlations as Probes of Dark Energy
- Conclusion

May detect neutrino mass with CMB lensing

- May detect neutrino mass with CMB lensing
- Probe r ~ 0.01 with AdvACT and r~0.001 with CMB-S4

- May detect neutrino mass with CMB lensing
- Probe r ~ 0.01 with AdvACT and r~0.001 with CMB-S4
- Explore Dark Energy with CMB in many ways including CMB halo lensing and CMB lens crosscorrelations with redshift and shear surveys

- May detect neutrino mass with CMB lensing
- Probe r ~ 0.01 with AdvACT and r~0.001 with CMB-S4
- Explore Dark Energy with CMB in many ways including CMB halo lensing and CMB lens crosscorrelations with redshift and shear surveys
- Future CMB surveys are important complement to large-scale structure surveys